



Life PollinAction



With the contribution of the Life Programme of the European Union LIFE19 NAT/IT/000848

Guidelines

CREATION AND MANAGEMENT
OF HABITATS FOR POLLINATORS

URBAN AREAS

COORDINATING BENEFICIARY:



Università
Ca'Foscari
Venezia

**Dipartimento di Scienze Ambientali,
Informatica e Statistica**

CA' FOSCARI UNIVERSITY OF VENICE

Department of Environmental Sciences, Informatics and Statistics

Via Torino 155

30172 Mestre (Venice) - ITALY

tel. 041 2347738 - 041 2347741

www.lifepollinaction.eu

lifepollinaction@unive.it



LIFE_POLLINATION



[lifepollinaction](https://twitter.com/lifepollinaction)

PROJECT PARTNERS:



Publication produced under LIFE19 project NAT/IT/000848.

Reproduction is authorised, citing the source.

Recommended citation: Buffa G., Baldin M., Fantinato E., Fiorentin R., Gastaldi A.M., Lorenzato L., Preo S., Scarton F., Toniolo C., Tosin M., 2024. Creation and management of habitats for pollinators. Guidelines. Project LIFE PollinAction (LIFE19 NAT/IT/000848). II. URBAN AREAS

Photos and illustrations are the property of LIFE PollinAction, unless otherwise stated.

Photos by:

E. Fantinato: page 29

L. De Savi: left side page 11

M. Valecic: page 22

R. Corrado: right side page 11, second from the right side page 11

R. Fiorentin: right side page 23, 36, 38

S. Fabian: left side page 23, 30, 31, 44, 50

S. Vaccher: second from the left side page 11

Graphic design and layout:

ALBATROS SRL - tel. 0461 984462 - info@albatros.tn.it

With the contribution of the LIFE Programme of the European Union.

Guidelines

CREATION AND MANAGEMENT
OF HABITATS FOR POLLINATORS

URBAN AREAS



THE LIFE POLLINATION PROJECT ACTIONS FOR BOOSTING POLLINATION IN RURAL AND URBAN AREAS LIFE19 NAT/IT/000848

THE PROJECT

The LIFE project PollinAction aims at the implementation of concrete actions to protect pollinating insects and to address the current ‘pollination crisis’.

The causes of this phenomenon are numerous, but the most substantial impact can be attributed to human land use. The processes of urbanisation and the expansion of intensive agricultural areas have resulted in a profound simplification of the landscape and the disappearance of habitats that were once widespread. Meadows, small woods, and hedges, as well as field and road edges, are becoming increasingly rare and reduced in size. Yet, it is precisely in these habitats that pollinators can find the resources they require to survive, including pollen and nectar, as well as areas in which they can nest and withstand the winter. In such simplified landscapes, ensuring the availability of habitats suited to the needs of pollinating insects is the only effective strategy to favour their presence and activity. Small areas in which these animals can find food resources, but also reproductive sites, and which allow them to safely move around the territory: meadows, small, wooded patches, wetlands, hedges, extensive agricultural areas. These interacting components create a network, a Green Infrastructure, capable of providing many benefits to both wild species, plants and pollinators, and humans.

WHO THEY ARE

The project, co-financed by the European Union, is co-ordinated by Ca’ Foscari University of Venice and developed in collaboration with the Autonomous Region of Friuli-Venezia Giulia - Central Directorate for Agri-food, Forestry and Fishery Resources, Biodiversity Service; Veneto Region - Agri-environmental, Fish and Wildlife Planning and Management Directorate; Veneto Agency for Innovation in the Primary Sector - Veneto Agricoltura; Municipality of Caldogno; Concessioni Autostradali Venete - CAV S.p.A.; Centro de Investigación y Tecnología Agroalimentaria de Aragón; ALBATROS S.r.l.; EcorNaturaSi S.p.A.; SELC soc. coop..

THE AREAS

Italy (Veneto and Friuli-Venezia Giulia) and Spain (Aragon). Three settings: agricultural areas, urban areas, and road infrastructures.

THE OBJECTIVES

LIFE PollinAction aims to mitigate the pollinator crisis by creating or improving their habitats, primarily by increasing the abundance of wild plants.

Specifically, LIFE PollinAction activities are dedicated to:

- creating approximately 28 hectares of new habitats rich in plant species, including 18 hectares of new meadows, 7 hectares of flowering strips, and 2 hectares of shrub patches;
- creation of 14 kilometres of new hedges;
- improvement of 230 hectares of meadows;
- assessment of ecosystem services provided by habitats and design and implementation of Payments for Ecosystem Services (PES) schemes to inform regional/national agricultural policy;
- designing milk, honey, and hay supply chains based on species-rich meadows to increase farmers' competitiveness.

TABLE OF CONTENTS

PREFACE	6
Why a guide?.....	7
Who is this guide for?.....	8
How to use these Guidelines.....	8
SECTION 1	9
What is pollination.....	10
Who are pollinators?.....	11
Why are pollinators important?.....	13
Plants and pollinators.....	13
Why are pollinators disappearing?.....	14
SECTION 2	15
What is the goal?.....	16
Not just pollinating insects!.....	16
How much space do I need?.....	17
What should I pay attention to?.....	21
Material selection.....	21
Method selection.....	28
Site condition.....	28
How many different species?.....	28
Design habitats to cover the entire flowering period.....	32
Management.....	32
SECTION 3	35
Meadow improvement.....	36
Meadow improvement: how to do it.....	36
Creation and management of wildflower strips.....	43
Creation of perennial flower strips: how to do it.....	44
Management and maintenance of wildflower strips.....	46
Creation and management of shrub patches and field hedges.....	49
The realisation of shrub patches and field hedges: how to do it.....	51
Management and maintenance of hedges and shrub patches.....	56
SECTION 4	58
Guide to assessing the quality of parks, gardens and courtyards.....	59
APPENDIX 1	81
Intervention Timeline.....	82

A photograph of a field of flowers, likely Delphinium, with purple and white blossoms. The word "PREFACE" is overlaid in the center in a white, bold, sans-serif font. The background is a clear blue sky and some blurred greenery.

PREFACE

WHY A GUIDE?

The expansion of urbanised areas is a continuous process, with the consequence that, at the present time, more than half of the world's population (approximately four billion people) live in urban areas.

The expansion of urban areas has significant and long-lasting impacts on plant and animal biodiversity, largely due to the reduction of green spaces, including natural and semi-natural habitats, which provide essential resources and refuge for wild species. When green spaces are present, they represent small, invaluable islands within an otherwise inhospitable matrix.

Urban areas can also serve as unusual habitats for wild species, particularly insects. Gardens, courtyards, and parks, for instance, can offer abundant and diverse resources, especially in comparison to those available in the agricultural areas surrounding the city. This is particularly true when these agricultural areas are subject to intensive agriculture and the frequent use of insecticides and pesticides.

The implementation of effective green space management strategies in urban and suburban settings can facilitate the provision of essential habitats, food sources, and protection for diverse plant and animal species, thereby contributing to the maintenance and enhancement of biodiversity. This, in turn, enables the transformation of our cities from potential threats to vital refuges for biodiversity.

The objective is ambitious, but integrating nature into urban environments through strategic urban growth planning and the designation of green spaces can yield numerous additional benefits. Green areas sequester carbon, attenuating the effects of climate change. They enhance local climates, reducing energy consumption. Urban green spaces improve urban aesthetics, stimulate active and healthy lifestyles, improve mental health, and provide venues for social interaction.

The indications provided in these Guidelines are a synthesis of the experience gained during the LIFE PollinAction project, with the aim of providing technical and practical information for environmental restoration projects in urban and peri-urban areas. In particular, the guide supports the European Union Pollinator Initiative¹, adopted by the Commission in 2018, by outlining the steps needed to plan, create, and maintain optimal natural and semi-natural habitats for pollinators.

This guide, which is dedicated to urban areas, forms part of a series that also includes a guide dedicated to the agricultural setting and another on road infrastructure.

¹ https://environment.ec.europa.eu/topics/nature-and-biodiversity/pollinators_en

WHO IS THIS GUIDE FOR?

This guide has been developed for municipal authorities and citizens who are interested in taking action to conserve or create habitats for pollinating insects on their land.

Municipalities are key actors: they are in a unique position to leverage their political influence and responsibility to advance the sustainable utilisation of urban green spaces and adopt strategic plans at the municipal level to dispose of land for the expansion or creation of new habitats for pollinating insects, involving local volunteers and Third Sector actors, especially in management activities.

Technicians and **Professionals** involved in landscape, housing and infrastructure planning may find it beneficial to possess a comprehensive understanding of the techniques employed in the creation, management and enhancement of habitats, as well as the goods and services they produce.

Private landowners and **Citizens** can contribute by creating or preserving habitats for pollinators on their land, which are essential for creating ecological connectivity within the area.

Voluntary associations and **Third Sector** actors can become players in the active management of urban green spaces, the real social capital of local communities.

HOW TO USE THESE GUIDELINES

Section 1 introduces the topic of pollinators and pollination.

Section 2 explains key strategies to support pollinators in urban areas.

Section 3 provides concrete indications for the improvement of habitats for pollinators.



Section 4 puts forth a straightforward questionnaire to evaluate the quality of parks, gardens, and courtyards and to ascertain straightforward measures to render them more 'insect-friendly'.

SECTION 1

WHAT IS POLLINATION?

Pollination can be defined as the transfer of pollen between flowers of different plants of the same species. It represents an essential phase of the reproductive cycle of the majority of flowering plants, including numerous species that provide us with food and materials. This process is facilitated by the movement of insects or other animals from one plant to another. In the absence of pollinators, many plants would be unable to produce seeds and reproduce, which would result in a decline in plant diversity and, subsequently, the loss of resources for insects.

POLLINATION



WHO ARE POLLINATORS?

It is perhaps unnecessary to provide an introduction to pollinators, as many are well-known. The honeybee is undoubtedly the most renowned pollinator, and its decline has attracted considerable attention from the general public. However, it is a common misconception that the honeybee is the most significant pollinator. In fact, the majority of pollination is performed by wild pollinators. It is estimated that there are at least 20,000 species of wild bees, in addition to bumblebees, butterflies, wasps and ants. These species are of vital importance for the preservation of the environment, both natural and cultivated. The principal pollinators can be classified into four main groups.

WHO ARE POLLINATORS

HYMENOPTERA

Bumblebees and bees, both social and solitary, wasps

Bees are among the most efficient pollinating insects in handling flowers, and many plants, both wild and cultivated, form obligate relationships with them. Wasps also contribute to the pollination of many plant species and, like bees, require pollen and nectar for their survival. In some cases, ants can also make an important contribution to pollination.



A specimen of *Bombus pascuorum* (Common Carder Bee)

LEPIDOPTERA

Butterflies and moths

They visit a wide variety of wild plants in search of nectar. They are less effective than bees in transporting pollen. Characterised by elongated, slender legs and a specialised tubular proboscis-shaped mouthpart, called spirotrumpet, they are not able to manage to intercept much pollen with their bodies nor do they have special structures to collect it.



A specimen of *Macroglossus stellatarum* (Hummingbird Hawkmoth) on *Scabiosa* sp.

DIPTERA

Flies, including hoverflies

They are an important yet often underestimated group of pollinators. Diptera are distinguished from other insects by their two membranous forewings and two reduced hindwings, which serve as balancing organs during flight. They are an ancient group and were probably among the first groups of pollinators. Diptera visit flowers to obtain nectar, which provides energy, and pollen, necessary for egg production.



A specimen of *Ectophasia crassipennis* (Thick-legged Ectophasia) on *Pimpinella saxifraga* (Burnet-saxifrage)

COLEOPTERA

Beetles and many others

They are a very ancient and incredibly diverse group of pollinators. More than 300,000 species have been described worldwide, representing approximately 40% of known insects. Beetles have contributed to shaping the earliest pollination relationships between plants and insects.



A specimen of *Cetonia aurata* (Rose chafer) on *Prunus mahaleb* (Mahaleb cherry)


BOX: WHAT THEY LOOK LIKE AND HOW THEY LIVE

Some insects do not have wings and flies have only one pair of functioning wings. However, in Coleoptera, the first pair of wings is modified to form a protective covering for the second pair of wings

The **insect body** consists of three parts:


- > the **head** on which are the eyes and antennae,
- > the **thorax** to which are articulated 3 pairs of legs and, in most cases, 2 pairs of wings,
- > the **abdomen**.

While we have bones inside our bodies, insects have an external skeleton (**exoskeleton**) that surrounds their bodies like a kind of shell.



Antennas are the nose, ears and tongue of insects and also host the sense of touch

Insects are **oviparous**, emerging from eggs. In some species the young insects are morphologically similar to adults (**heterometabolous insects**). In many other species, development occurs through several stages, each different in appearance, shape and size: larva, pupa, perfect insect or adult (**holometabolous insects**).



The lifespan of insects exhibits considerable variation between species, with some species having a lifespan of just a few days and others reaching several years.

WHY ARE POLLINATORS IMPORTANT?

Pollinators are of critical importance in maintaining healthy ecosystems and ensuring food production, as they facilitate the reproduction of flowering plants. The role of pollinators in the functioning of our agricultural system is of fundamental importance. It is estimated that approximately 90% of flowering plants and 75% of food crops worldwide are dependent on pollinators.

It is possible to consider the use of artificial pollination or the purchase of bees and bumblebees as potential solutions to the problem. But without wild plants? Pollinators are crucial for the reproduction of wild plants. Without them, plant populations would decline, even if other essential elements, such as soil, air, and nutrients, were available.

PLANTS AND POLLINATORS

Insect pollination represents one of the most remarkable examples of co-evolution, whereby two distinct species, a plant and an insect, engage in a symbiotic relationship that is mutually beneficial, resulting in evolutionary changes for both parties. The relationships established between these two groups of organisms give rise to the high degree of complexity and diversity observed in nature.

Flowering plants are amazing! They adapt to their pollinators by developing structures and strategies to attract specific pollinators. Insect-pollinated flowers often exhibit bright colours and patterns, which are often invisible to the human eye, that guide insects towards pollen and nectar. And it is a two-way street! Insects adapt to plants by evolving body parts and behaviour to favour the pollination of specific plants. Bees, for example, have developed special sucking mouthparts to absorb nectar, and in some species, they even have small baskets on their hind legs to collect pollen.

About 90% of plants require the help of other organisms, such as insects, birds, or bats, to transfer pollen from one plant to another. Thanks to the pollination process, seed plants can produce seeds and fruit; likewise, plants provide insects with the resources, pollen and nectar, as well as reproductive and overwintering sites necessary to complete the life cycle.

WHY ARE POLLINATORS DISAPPEARING?

The decline of pollinators is the result of a multitude of interrelated factors, including the use of pesticides, the spread of diseases and pests, climate change, and the introduction of alien species. However, the most significant impact can be attributed to human land use practices. The expansion of urbanisation and the intensification of agricultural management have contributed to the simplification of the landscape and the reduction, and in some cases, the loss of natural and semi-natural habitats that are vital for pollinators.



The process of urbanisation has a profound and enduring impact on the biodiversity of plant and animal species. This is largely due to the reduction of green spaces, which, when present, represent small hospitable islands within an otherwise inhospitable urban matrix

SECTION 2

WHAT IS THE GOAL?

Pollinators are adversely impacted by a multitude of factors, with the loss of natural and semi-natural habitats, such as meadows, hedges, and woodlands, representing a significant concern. The loss of habitat is accompanied by a decline in the diversity and abundance of plant species that provide essential resources for pollinators, including food (pollen and nectar) and shelter.

The recommendations set forth in these Guidelines are intended to **address the decline in biodiversity and enhance the ecological integrity of urban ecosystems**. This is achieved by creating or enhancing habitats that are rich in entomophilous flowering species², which can provide nutritional resources (pollen and nectar) and ensure the availability of reproductive sites for pollinating insect populations. The presence of these elements within urban environments can lead to a significant increase in the richness and abundance of insects.

NOT JUST POLLINATING INSECTS!

The establishment of hedges, meadows and wildflower strips in schoolyards, parks and gardens, flowerbeds and roadsides provides an effective refuge for a multitude of plant species and a variety of animal species, including birds and other wildlife.

The creation and maintenance of habitats and the flora that inhabit them offers the opportunity to achieve a variety of benefits through a single action. These include the regulation of temperature and the mitigation of the heat island phenomenon in urban centres, the reduction of surface water runoff in the event of increasingly frequent intense precipitation, and the provision of opportunities for outdoor exercise and recreation. These are just a few of the benefits that green areas in urban environments provide. The presence of green areas has a relaxing and restorative effect, and contributes to the aesthetic appeal of cities, thereby facilitating the maintenance of a harmonious relationship between humanity and the natural environment.

² *Plants pollinated by insects, especially bees, butterflies, moths, dipterans and beetles*

BRINGING NATURE BACK INTO OUR CITIES PROVIDES US WITH MANY BENEFITS

Health and psycho-physical well-being
urban green spaces offer important recreational and wellness benefits to residents

Scenic quality
urban green spaces enhance aesthetic quality

Pollination
habitats and resources for pollinating insects



Biodiversity
species, habitats

Heat wave mitigation
urban green spaces have a great capacity for cooling and mitigating temperatures

Runoff control
control of surface runoff by green spaces can prevent flooding, reducing damage to population and infrastructure

The main benefits provided by natural and semi-natural habitats in the urban environment. Yellow boxes indicate services directly related to the overall functioning of ecosystems; grey boxes indicate services related to society and the environment.

HOW MUCH SPACE DO I NEED?

Schoolyards, parks, backyard and vegetable gardens, flowerbeds, roundabouts, and roadsides all have the potential to meet the minimum needs of pollinators. This means that they can provide the resources (pollen and nectar) that are essential for the insects to survive, as well as the opportunity for them to nest and survive the bad season.

Size is not always important. The variety and quality of the habitats created or improved are more important than quantity! It is possible to place plants in a variety of settings: small or large parks and gardens, but also in pots and on windowsills. Planting a wide variety of flowers, with different colours, flower sizes and flowering times (spring to autumn), will attract a wide range of pollinators and ensure a continuous supply of resources for both adults and larvae.

All city parks, and even private gardens contain small, little-used marginal areas that are just waiting to be discovered! Think of the small grassy patches and areas around buildings, and even roadsides, traffic islands and roundabouts, and the edges of drainage channels, can be used for this purpose. Even small bare patches are crucial for all those insects that nest in the soil. Despite the vast diversity of pollinator species, they all require three types of habitats to survive and complete their life cycle:

- **foraging habitat:** areas characterised by the presence of floral resources (pollen and nectar), within a reasonable flying distance, spanning from spring to late autumn. The greatest number of pollinators and, consequently, the greatest demand for resources occurs in early summer. However, the availability of floral resources must also be ensured in spring, when overwintering pollinators emerge from hibernation and require sustenance, and in late summer, when pollinators need to feed in preparation for hibernation;
- **breeding habitat:** breeding habitats are species-specific. For example, some bee species (both solitary and social) construct their nests in bare soil, either by digging or by utilising abandoned nests and burrows. Other bee species utilise leaves, while others employ cavities in plant stems or walls. Butterflies require specific native plants, both herbaceous and shrubby species, which serve as a food source for their caterpillars;
- **overwintering habitat:** areas with vegetation that remain undisturbed from late summer until late spring, such as permanent meadows, road or ditch margins, shrub patches, hedges, groves, and isolated trees, as well as timber piles and litter.



Pontia edusa on *Sanguisorba*

WILD POLLINATORS: FOOD REQUIREMENTS AND HABITAT

POLLINATORS	DIETARY REQUIREMENTS	BREEDING HABITAT	OVERWINTERING HABITAT
Solitary bees <i>(Hymenoptera)</i>	Pollen and nectar	Most of the species nest in bare or partially vegetated, well-drained soil. Others nest in narrow tunnels in the trunks of dead trees. Some species build their nests by folding leaves over themselves.	Stems of dead plants, bark, reeds, foliage, and especially undisturbed soil are the winter homes of solitary bees.
Bumblebees <i>(Hymenoptera)</i>	Pollen and nectar	The majority nest in small cavities, often underground, in abandoned rodent nests, or in tree hollows, at the base of tufts of grass, or in wall crevices.	The queens overwinter underground in shaded areas, usually near trees, on embankments, and on north-facing slopes, to avoid emerging too early on a warm or sunny day in winter.
Wasps <i>(Hymenoptera)</i>	Pollen and nectar	Most wasp nests are made of paper. Wasps utilise saliva and chewed wood pulp to create their nests. The appearance of the nests varies depending on the species that built them. They are typically located in sheltered areas with easy access to the outside, such as hollow trees, tall grass, or buildings. Other species build their nests of mud in sheltered locations.	Some adult wasps overwinter in protected and undisturbed locations, such as under tree bark or in building cavities.
Butterflies <i>(Lepidoptera)</i>	Nectar	Butterflies construct their nests in proximity to plants that are utilised by caterpillars as a source of nourishment.	Most butterflies survive the winter as a chrysalis or larva (caterpillar) in vegetation or litter; others overwinter as adults in sheltered cavities or tree trunks; still others migrate over long distances.
Hoverflies <i>(Diptera)</i>	Nectar and pollen	The females lay eggs in suitable places where the larvae can find food. Species that feed on aphids lay eggs near aphid colonies, species with larvae that feed on plants lay eggs on host plants, while other species lay eggs in stagnant water.	Hoverflies have two strategies to survive the winter. They either burrow into the ground as larvae or enter a state of hibernation as adults in sheltered places within the trunks of old trees or in building crevices.
Beetles <i>(Coleoptera)</i>	Pollen	Some species lay eggs among decomposing leaves, others in decaying wood, or in animal feces.	Adults survive the winter in tree cavities, among decomposing leaves, under logs, and in the soil.

BOX: FLOWERS FOR FREE

One of the most effective strategies for the conservation of pollinators is the maintenance and enhancement of existing habitats. Some species of plants that are important for pollinators are very common in parks, backyards, urban gardens or roadsides.

SPECIES	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
<i>Achillea millefolium</i>												
<i>Buphthalmum salicifolium</i>												
<i>Calystegia sepium</i> subsp. <i>sepium</i>												
<i>Centaurea nigrescens</i>												
<i>Cichorium intybus</i>												
<i>Cirsium vulgare</i>												
<i>Crepis biennis</i>												
<i>Daucus carota</i> subsp. <i>carota</i>												
<i>Geranium molle</i>												
<i>Glechoma hederacea</i>												
<i>Helminthotheca echioides</i>												
<i>Lamium purpureum</i>												
<i>Leontodon hispidus</i>												
<i>Iris pseudacorus</i>												
<i>Lotus comiculatus</i>												
<i>Lythrum salicaria</i>												
<i>Ranunculus acris</i>												
<i>Ranunculus bulbosus</i>												
<i>Salvia pratensis</i>												
<i>Taraxacum officinale</i>												
<i>Torilis arvensis</i>												
<i>Trifolium pratense</i>												
<i>Trifolium repens</i>												
<i>Veronica arvensis</i>												
<i>Vicia sativa</i>												

LEGEND: The coloured squares represent the colour of the flowers and the flowering period. Plants to the left of the vertical dotted line are of particular importance in providing floral resources for insects during the early stages of the season.

WHAT SHOULD I PAY ATTENTION TO?

MATERIAL SELECTION

The selection of the appropriate material, the plants, to be used is of paramount importance to ensure the success of the intervention, reduce the cost of cultivation treatments, and at the same time maintain the integrity and stability of natural ecosystems. It is imperative that the plants be of '**local provenance**', whereby the seed used for plant production must have been collected in the same 'region of provenance' (ROP) as that of the plants' destination. In the event of unavailability, it may be accepted that the material sourced from other ROPs may be used, provided that it is sourced from locations as close as possible to the site of intervention.

In the case where interventions entail the utilisation of **seedlings in soil blocks**, at the time of delivery, these must be provided with soil firmly attached to the roots and free from weed plants; they must be free from obvious pathologies, with sufficient development of the above-ground part to facilitate effective handling by operators. Woody plants, trees and shrubs, must be well lignified, preferably

BOX: DEFINITION OF REGION OF PROVENANCE

For plant species, the region of provenance is defined as 'the territory or set of territories subject to sufficiently uniform ecological conditions and on which there are topsoil or seed sources that are sufficiently homogeneous from a phenotypic and, if evaluated, genotypic point of view, taking into account altitudinal limits where appropriate' (Directive 1999/105/EC on the marketing of forest reproductive material). For forest species, regions of provenance are defined nationally by Ministerial Decree No. 9403879 of 30/12/2020 'Establishment of the National Register of Basic Materials' and subsequent amendments and integrations, and possibly detailed by regional regulations. The recent Ministerial Decree No. 269708 of 11/06/2021 'Division of the Italian territory into Regions of Provenance' includes a cartographic representation of the aforementioned regions of provenance. This tool, available online³, can also be adopted as a precautionary measure for herbaceous species, marketed in various forms (seedlings, seeds, brush-harvested seed, bulbs, etc.) and for which no analogous legislation exists. An additional classification⁴ is also available online, comparable to the previous one (although based on different criteria) and equally suitable when adopting a precautionary approach in the handling of plants and seeds. The provenance of plant material is typically guaranteed by public nursery facilities that, for institutional purposes, produce only locally sourced plants. In any case, the customer is required to verify the information regarding the origin of the plants, requesting documentation from the supplying nursery (which may or may not coincide with the producing nursery), as suggested below:

- if the plant nursery producing the plants is certified according to UNI EN ISO 22005 (traceability), it is sufficient for the nursery to declare, before delivery, the denomination (location and municipality) of the stand from which the batches supplied were collected;
- if the producing nursery is not certified according to traceability, the customer is expected to request the following information: the source population (location, municipality) of the seed used, the year of collection of the same, the quantity used for the production of that batch of plants, and the corresponding number of plants obtained.

³ <https://www.politicheagricole.it/flex/cm/pages/ServeBLOB.php/L/IT/IDPagina/17155>

⁴ *Classification and mapping of the ecoregions of Italy.*

<https://www.tandfonline.com/doi/full/10.1080/11263504.2014.985756>

of an age less than or equal to two years (with a maximum age of three years), with a maximum height of 80–90 cm and a minimum height of 20 cm. In instances where the root system has become excessively twisted and compact due to prolonged nursery cultivation, it may be beneficial to sever the root ball at the end prior to planting. This can assist in the disentanglement of the roots and facilitate the establishment of a robust root system, thereby enhancing the speed of the seedling's establishment.



If the top of the root system appears twisted, before planting, it may be useful to sever the end of the soil block to facilitate the disarticulation of the roots and accelerate the establishment of the young seedlings

BOX: WHY NATIVE PLANTS

The term 'native plants' (also referred to as 'autochthonous') denotes species that have originated and evolved within the geographical area in which they are found. Consequently, they have a long history of natural presence within the natural ecosystems of a given region. Non-native plants, also referred to as allochthonous, alien, or exotic plants (*for further information, please see the BOX Exotic Species on page 24*), are naturally widespread in other parts of the world, or even in other areas of the country (for example, they are typical of Mediterranean climate areas). However, they can also grow well in other areas as ornamental or garden plants. The use of native plants, which are adapted to the local climate and conditions, is a more effective method of ensuring the success of the intervention and its long-term duration. It is of the utmost importance to note that, although exotic plants may offer pollinators some nutritional value, native plants, due to their long history of co-evolution, attract and support a greater diversity of native pollinators and are crucial for the survival of numerous species.

LIFE PollinAction's Public Reference Nurseries

The plants used for the creation and/or improvement of habitats in the LIFE PollinAction Project were produced and supplied by public nurseries, project partners.

- Centro Biodiversità Vegetale e fuori Foresta di Veneto Agricoltura a Montecchio Precalcino (VI): Via Bonin Longare, 6, 36030 - tel. +39 0445 864445;
- Regional nursery centres of Friuli-Venezia Giulia:
 - > Azienda Volpares a Palazzolo dello Stella (UD): Casali Volpares, near Palazzolo dello Stella (UD);
 - > Vivaio forestale Pascual di Tarcento (UD): Via Pradandons, 15 - tel. +39 0432 785029.



Cultivation of seedlings in soil blocks at Volpares Nursery (Palazzo dello Stella)



Seeds of native species of hay meadows stored at Veneto Agricoltura Nurseries (Montecchio Precalcino)



Plots for intensive production of local seed for hay meadows at Veneto Agricoltura Nurseries (Crespano del Grappa)



BOX: EXOTIC SPECIES

A significant proportion of the plant species utilised in agricultural, horticultural and forestry contexts are not native to our country. Rather, they have been introduced from other regions of the globe. Such species are designated as 'exotic' or 'alien'. The majority of these introductions have resulted in benefits, as evidenced by the significant role of exotic species in the agri-food sector. However, some of these introduced species have become naturalised in the regions of introduction and subsequently spread uncontrollably, invading both natural ecosystems and agricultural and urban environments. This has resulted in a range of negative impacts. A typical example of an invasive species along roadsides is Johnson grass (*Sorghum halepense*). These particular exotic species are defined as invasive alien species because they have the capacity to alter, and in some cases, irreversibly disrupt the structure and functioning of ecosystems, with adverse ecological and economic consequences. In some instances, these species have also been shown to have detrimental effects on human health.

Invasive species exhibit a number of characteristics: the alarm bells!

- ability to colonise disturbed environments and bare ground
- very rapid growth
- short life cycle with early flowering and dissemination, often different from that of native species
- production of large quantities of seeds and/or fruits
- effective propagation through vegetative means

It is these very characteristics that render these species so popular for use as ornamental greenery.

However, the introduction of invasive exotic species represents a significant threat to biodiversity. It is therefore imperative to adopt responsible practices, including the avoidance of invasive plant species, to prevent further disruption to natural ecosystems.

The phenomenon of invasions is experiencing a marked increase, with a considerable number of new species being introduced on a voluntary or accidental basis each year. In view of this situation, the European Commission has adopted a list of invasive alien species of EU relevance, which is updated on a periodic basis in accordance with EU Regulation 1143/2014⁵. This list identifies those invasive species whose negative effects are so significant that they require coordinated and uniform action at the EU level.

⁵ <https://eur-lex.europa.eu/legal-content/IT/TXT/?uri=celex%3A32014R1143>

SPECIES TO AVOID

TREES

Box elder Maple (*Acer negundo*)

Tree of Heaven (*Ailanthus altissima*)

Cigartree (*Catalpa bignonioides*)

Chinese windmill palm (*Trachycarpus fortunei*)

Black Locust (*Robinia pseudoacacia*)

SHRUBS

Butterfly Bush (*Buddleja davidii*)

False Indigo bush (*Amorpha fruticosa*)

Cherry Laurel (*Prunus laurocerasus*)

Oval-leaved Privet (*Ligustrum ovalifolium*)

Chinese Privet (*Ligustrum sinense*)

Glossy Privet (*Ligustrum lucidum*)

Oregon Grape (*Mahonia aquifolium*)

Japanese Rose (*Rosa rugosa*)

Japanese Spirea (*Spiraea japonica*)

HERBACEOUS PLANTS

Panicled Aster (*Symphyotrichum lanceolatum*)

Balsam (*Impatiens* spp., various species)

Japanese Honeysuckle (*Lonicera japonica*)

Evening Primrose (*Oenothera* spp., various species)

Phacelia (*Phacelia tanacetifolia*)

Tawny Daylily (*Hemerocallis fulva*)

Buckwheat (*Fagopyrum esculentum*)

Canada Goldenrod (*Solidago canadensis*)

Giant Goldenrod (*Solidago gigantea*)

Japanese Knotweed (*Fallopia japonica*)

Kudzu (*Pueraria lobata*)

American Pokeweed (*Phytolacca americana*)

The main introduction pathway of alien plant species into regions beyond their native range is through trade and use as ornamental plants. Indeed, approximately 80% of the invasive species currently present in Europe have been introduced voluntarily as ornamental plants. These are the plants we often find in private gardens or public green spaces.









One of the primary factors motivating managers or private citizens to select an ornamental plant is its aesthetic appeal, regardless of its provenance. The aesthetic appeal of a plant can be quantified through a series of morphological measurements, including the size of the plant, the dimensions of its flowers, and the hue of their colour. Once these parameters have been collated, a comparison can be made between two plant species in order to ascertain their aesthetic similarity. This method may prove useful in suggesting local (native) plant species in place of invasive alien species that better meet the aesthetic requirements of the green project to be realised.



A comparable analysis was undertaken to evaluate the similarity between the local flora and the ornamental species most commonly sold by nurseries and florists in the province of Venice. Of the 75 alien ornamental plant species analysed, nine were found to have at least one local species that was 100% aesthetically similar according to the chosen parameters. In addition, 42 other plants exhibited at least one local species with a similarity rating of 83%.

The utilisation of plant species native to the local flora in public and private green projects offers a multitude of advantages:

- reinforces the gene pool of the local flora by reducing the risk of hybridisation;
- increases the abundance and diversity of pollinating insects;
- facilitates the creation of more suitable habitats for the support of local bird populations;
- enhances the sense of belonging and reinforces the connection between individuals and the territory;
- can represent a novel market opportunity for local companies;
- particularly, thermophilous plant species are well adapted to urban environments, in which they find characteristics similar to the Mediterranean environment: limited water availability, high temperatures and brightness.

ALIEN ORNAMENTAL SPECIES	NATIVE SPECIES	ALIEN ORNAMENTAL SPECIES	NATIVE SPECIE
 <p><i>Camellia japonica</i></p>	 <p><i>Cistus creticus</i></p>	 <p><i>Primula japonica</i></p>	 <p><i>Primula farinosa</i></p>
 <p><i>Dimorphoteca pluvialis</i></p>	 <p><i>Tragopogon porrifolius</i></p>	 <p><i>Rosa gr. 'Pink-Favorite'</i></p>	 <p><i>Rosa gallica</i></p>

Code of conduct: replacing ornamental plants with native species

In 2018, the LIFE ASAP project produced a Code of Conduct for Nurseries and Invasive Alien Species. The Code of Conduct comprises 12 fundamental principles, one of which encourages nurseries and garden centres to make available replacement species to invasive alien species, possibly belonging to the local flora.

To learn more:
www.minambiente.it/pagina/specie-esotiche-invasive
www.specieinvasive.it
www.lifeasap.eu

METHOD SELECTION

As will be seen in Section 3, there are several techniques that can be employed for the creation or enhancement of habitats. The selection of the optimal method is contingent upon a number of variables, including the surface area to be treated, the practitioner's experience and technical preparation, and the availability of specific equipment. However, the methods are not mutually exclusive, and often, the best result is achieved by integrating various techniques. In all cases, it is essential to consider the **availability of materials**. It is important to note that not all species are part of the typical nursery production, and thus it is essential to plan the intervention well in advance.

SITE CONDITIONS





The growth of plants is subject to several environmental factors, which exert a significant influence on their development. Each plant species has a defined tolerance range in relation to the various ecological factors (light, temperature, humidity and soil characteristics) within which it is able to perform its essential functions and complete its life cycle. To ensure the success of interventions, it is of the utmost importance to pay particular attention to the plant mix, which must be designed in such a way so as to utilise species that are appropriate to the intervention site.

HOW MANY DIFFERENT SPECIES?

More is better! The richer and more diversified a habitat is, the more able it is to withstand a wide variety of environmental conditions. More importantly, the richer a habitat is in terms of plant species, the greater the number of pollinators it will attract. A diversified habitat ensures the presence of species with different floral traits, capable of satisfying pollinator insects with diverse feeding requirements and capabilities to manipulate flowers. Differences in floral traits such as flower shape, accessibility of resources, colour, etc., make different plant species suitable for different pollinator insects. For example, open flowers, with exposed pollen and nectar, such as those of the buttercup (*Ranunculus* spp.), are easily accessible to numerous different groups of pollinators, while flowers with more complex corollas, such as those of the common sage (*Salvia pratensis*), with pollen and nectar difficult to access, can be pollinated only by a narrower group of insects.

A high species richness allows for staggered flowering, i.e., different flowering periods that ensure the supply of resources throughout the season. Finally, increasing the number of different species ensures the presence of plants with different vegetative characteristics (leaves and stem), thus providing resources and breeding and overwintering habitats for different species of insects, including beneficial insects.

MAIN FLOWER TRAITS OF THE SPECIES VISITED BY DIFFERENT GROUPS OF POLLINATORS

POLLINATORS	COROLLA SHAPE	COLOUR	FRAGRANCE	MAIN FLORAL RESOURCE	EASE OF ACCESS TO FLORAL RESOURCES	EXAMPLES
HYMENOPTERA	Bilabiate	Blue and yellow	Sweet	Nectar	Limited	 <i>Ajuga reptans</i> , <i>Salvia pratensis</i>
LEPIDOPTERA	Tubular	Red and purple	Sweet	Nectar	Limited	 <i>Scabiosa triandra</i> , <i>Succisa pratensis</i>
DIPTERA	Disc	Yellow and white	Soft	Pollen	Good	 <i>Achillea millefolium</i> , <i>Pastinaca sativa</i>
COLEOPTERA	Disc	White and cream	Unpleasant	Pollen	High	 <i>Crataegus monogyna</i> , <i>Viburnum lantana</i>

1. Flowers with bilabiate morphology of *Lamium orvala* (Balm-leaved Red Deadnettle) - 2. Flowers with tubular morphology of *Lythrum salicaria* (Purple loosestrife) with pollinator (*Pieris rapae* - Small white butterfly) - 3. Head of *Knaulia arvensis* (Common ambretta) with pollinator (*Eristalis tenax* - Drone fly) - 4. Inflorescence of disk flowers of *Filipendula ulmaria* (Common Meadowsweet)

BOX: EVEN THE EYE WANTS ITS PART!

Pollinators do not only use size as a guide. They also use scents and colours. When using seedlings in soil blocks, it is recommended to implement a planting scheme that involves the formation of **small, mono-specific groups**, defined as small groups of plants of the same species. This method ensures that the plants are more visible and therefore more attractive to insects during the flowering period. Furthermore, since pollinators tend to visit neighbouring flowers, mono-specific groups facilitate the transfer of pollen between individuals of the same species. A useful tip is to alternate mono-specific groups of species with different flower structures.



Explanatory scheme of small mono-specific groups for the construction of wildflower strips

The specifics of planting schemes can vary depending on the needs.


In **wildflower strips**, mono-specific groups may comprise a variable number of plants, ranging from 4 to 10 plants. It is recommended that plants be spaced approximately 30-40 cm apart, but the distance should be assessed considering the final size of the plants and their propensity for vegetative propagation. For instance, for hygrophilous species like Water flag (*Iris pseudacorus*), which spread rapidly thanks to underground rhizomes, lower densities should be considered, such as 3 plants/m linear of the bank.

In the case of **meadows**, it is recommended to identify areas (plots) of about 2 sqm within which to carry out close planting of seedlings. The number of individuals per plot varies from 24 to 40, depending on the requirements and the size of the species at maturity. The minimum density of planting plots is 20 per 1,000 sqm, also variable depending on the needs.



Geometric layout for the construction of a flower strip

The arrangement of the plots may be random within the meadow or follow a straight line (transect), with a distance of approximately 3 m between each plot. In cases where the intervention involves rare or threatened species, lower densities than those indicated may be employed, yet still ensuring the creation of small populations capable of flowering, ripening and producing seed.



Explanatory scheme for planting species in plots to enrich species-poor meadows



Detail of the plots. The actual number of individuals should be assessed taking into account the final size of the plants and their propensity for vegetative propagation

DESIGN HABITATS TO COVER THE ENTIRE FLOWERING PERIOD

Ideally, a habitat should comprise a mix of species so as to ensure the provision of resources throughout the entire growing season, from April/May to September/October. Nevertheless, this is not always straightforward to accomplish.

The best strategy is to combine different habitats. For instance, many very common shrub species such as the Blackthorn (*Prunus spinosa*) and the Common hawthorn (*Crataegus monogyna*), exhibit very early flowering, thereby providing resources when herbaceous plants are not yet in bloom. The combination of different habitats enables the completion of the life cycle of different species. For example, many butterflies utilise meadows as a foraging habitat, while they require shrubs for egg-laying and nutrition during the larval stage. A typical combination can therefore be a hedge or a patch of shrubs, which provide resources in early spring and nesting and overwintering sites, and meadows and wildflower strips, which provide resources at the peak of the season, in summer and autumn.

MANAGEMENT

Proper habitat management is essential for their long-term maintenance. All habitats require some form of regular maintenance, both to ensure their initial establishment and to maintain them over time.

For **woody habitats** such as hedges and shrub patches, maintenance is required particularly during the **initial stages**, and especially the first year, to control invasive herbaceous species, which can otherwise have a detrimental impact on the growth and survival of young woody seedlings.

Conversely, **herbaceous habitats**, such as meadows and wildflower strips, need **regular annual maintenance** through mowing. The absence of periodic mowing gives rise to spontaneous dynamics processes within the habitat. Initially, the accumulation of dead plant matter results in a reduction in species diversity, particularly those that are entomophilous. Subsequently, the areas are colonised by brambles and shrubs, which are then succeeded by tree species. The practice of mowing serves to control the growth of weeds, maintain the diversity of plant species, and prevent the invasion of woody plants.

However, in urban environments, the primary challenges originate from intensive management practices, with regard to both the frequency of mowing and the height at which the grass is cut.

Frequent and close to the ground mowing results in a reduction in species richness. This is due to the fact that such mowing practices favour species that, due to their growth form, are

particularly well adapted to the disturbance caused by mowing, namely tufted grasses and annuals, as well as all plants that grow close to the ground, such as the common dandelion or clovers. In contrast, all species with an erect habit, such as daisies, bluebells and cornflowers, are progressively reduced. The change in plant species composition also has a cascading effect on the insect community, resulting in the loss of the benefits associated with high insect diversity, including pollination, biological pest control, and the decomposition of organic matter. In summary, although it may appear counterintuitive, **the practice of frequent mowing, undertaken primarily for aesthetic purposes, can give rise to other aesthetic issues as a consequence of its role in facilitating increased invasion of weeds and pests.**

It is equally important to consider the height of the cut. A meadow that is maintained at a higher height is more resistant to pests, weeds, and drought events. The optimal maintenance strategy is to limit mowing to a maximum of two or three times per year, with a minimum cutting height of approximately 20 cm. This should be conducted after the entomophilous species have flowered and dispersed, as this ensures their long-term persistence.

BOX: THE COST OF INTENSIVE MANAGEMENT

In addition to the change in plant species composition and the cascading effects this has on the entire community, it is worth mentioning that intensively managed meadows also require considerable investment in maintenance costs, as well as the utilisation of environmentally harmful fertilisers and pesticides.

In a recent study conducted in Canada⁶, the authors presented the financial implications of municipal meadow management, with costs calculated on a per-unit area basis (in hectares). The estimated cost for mowing 15 times a year was found to be 130 euros per hectare. The costs included the remuneration of operators, the costs of equipment and fuel, but did not consider the costs of pesticides or fertilisers.

A reduction in mowing frequency from 15 to 10 times per year for the same area resulted in a 36% reduction in costs.

While the mowing costs per hectare reported in the research may differ from those observed in other regions and jurisdictions, the positive economic impact of the management change is nevertheless considerable and worthy of note.

⁶ <https://besjournals.onlinelibrary.wiley.com/doi/epdf/10.1111/1365-2664.13542>

BOX: BEST PRACTICES FOR MOWING HABITATS FOR POLLINATORS

Mow no more than 2(3) times a year

An increase in the frequency of mowing can result in a reduction in species richness and diversity, with a notable decline in the number of species that are beneficial to pollinating insects

In case the area is subject to high levels of visitation, resulting in the necessity for more frequent mowing, it is recommended that no more than 50% of the area is mowed at any one time, or that uncut strips are maintained

This approach will facilitate the recolonisation of the mowed site by both plants and insects

Schedule mowing over time

Mowing should be scheduled after the entomophilous species have reached flowering and disseminated

In the event of colonisation by weed species, it is recommended that mowing be scheduled prior to the maturation of the seeds

The most cost-effective and successful method to eradicate weeds is to prevent them from producing and dispersing seeds

Use a minimum cutting height of 20-25 cm

Mowing at this height effectively controls invasive plants while minimizing the impact on native plants and many insects

Avoid the use of fertilisers

The application of fertilisers tends to favour a few highly competitive species, which in turn reduces the floristic richness of the meadow

SECTION 3

MEADOW IMPROVEMENT

The objective of this sheet is to provide a set of guidelines for the **enhancement of existing meadows that have undergone a decline in species diversity.**

In urban areas, the loss of species is predominantly attributable to intensive management practices, although a decline in management activity may also be a contributing factor.



MEADOW IMPROVEMENT: HOW TO DO IT

1. Recovery by re-establishing management

As evidenced in Section 2, in the chapter on Management on page 32, inadequate management practices result in a decline in species diversity, particularly entomophilous species, and the introduction of synanthropic-ruderal⁷ or exotic species that can be detrimental to the ecosystem. In such instances, the enhancement of meadow habitats can be achieved through the implementation of **well-designed management strategies.**

If meadow degradation is due to intensive management, the recovery of species richness can be achieved by decreasing the number of cuts and the cessation of any fertilisation practices (see *BOX BEST PRACTICES for mowing habitats for pollinators* on page 34).

⁷ *Synanthropic-ruderal species are typically small-sized plants that spread in habitats altered and disturbed by humans, such as roadside, urban areas, or abandoned cultivated fields.*

Conversely, if the impoverishment of the meadow is caused by a slowdown in management practices, the restoration of species richness can be achieved by implementing appropriate management strategies, which involves 2(3) cuts during each growing season, with removal of mowed material. The first mowing should be undertaken prior to seed dispersal by ruderal and exotic species, while simultaneously allowing for the maturation and seed drop of spring-summer native species, thus in the first fortnight of June. The second mowing, which is typically conducted from mid-August onwards, is undertaken with the objective of containing the proliferation of late-summer weed species (e.g., Johnson grass, *Sorghum halepense*). The aim of double mowing and hay removal is to deplete the soil of excess biomass and plant residues, in order to avoid new inputs of organic matter and accelerate settlement and balance over time. The objective is to bring the soil and meadow back to their original conditions, which are those most complex and rich in plant and animal biodiversity. In case the slowdown in management practices has resulted in the establishment of woody species, it is imperative to implement a programme of mechanical clearing, using a field shredder or brush cutter. The optimal timeframe for the initial mechanical shredding is late autumn or winter, when the plants are dormant and to avoid disturbing wildlife populations. If the areas are colonised by **brambles** or other **invasive exotic species** (e.g., *Ailanthus altissima* or *Amorpha fruticosa*), it is necessary to bear in mind that, due to their great capacity for propagation, the complete eradication of these species may require several passes, at each vegetative regrowth, to remove young shoots.

2. Recovery through species enrichment

If the improper management has been in place for an extended period of time, or in instances where the meadow is experiencing a notable decline in biodiversity, it may be feasible to pursue species enrichment through the implementation of diverse methodologies.

2a. Enrichment through overseeding

Overseeding is a technique used to regenerate and improve meadows. It involves the sowing of seeds on existing grassland characterised by low species diversity, particularly of entomophilous species.

It is of the utmost importance to ensure that the seed reaches the ground below the existing turf. Therefore, prior to sowing, the grass must be mowed with a very low cut with subsequent removal of the mowing.

Overall, methods are contingent upon the size of the area to be overseeded and the available equipment. For **large surfaces**, there are specific seed drills with a large seed hopper, coupled with a harrow to prepare the soil; a corrugated roller is employed to enhance seed penetration. Alternatively, the deployment of scarifiers/aerators can be considered. For **smaller areas**, a variety of small equipment is available on the market, including items such as aerators, rollers and manual seeders. At the hobbyist level, even a simple preliminary work with a rake and broadcasting seeding can be effective. However, it is essential that the seeds reach the soil and do not remain on the grass. In the case of overseeding, the quantity indicated in the literature as sufficient is 0.5-1 g/sqm.

2b. Enrichment through planting of native seedlings in soil blocks

This type of intervention involves the use of seedlings in soil blocks, i.e., pre-grown plant material.

The seedlings serve as ‘seed carriers’: once established, they can produce their own blooms and viable seeds for the spontaneous spread of species in the surrounding areas. This type of intervention enhances the quality of the habitat and its suitability for pollinators. The same approach can also be used to accelerate meadow enrichment with species of conservation interest, as they may be rare or threatened.



Native seedlings in soil blocks

BOX: WHY USE SEEDLINGS IN SOIL BLOCKS?

Compared to seeds, seedlings in soil blocks have the advantage of faster development, enabling them to rapidly compete with existing species, thus avoiding strong competition during the initial growth phases, which are particularly difficult and delicate for plants, occurring in open field conditions.

The **optimum time for planting is autumn**, as for all other operations involving plant material. Seedlings in soil blocks should be planted in the soil after making a hole suitable for the size of the underground part (root ball or soil block). The plant should be immersed in the soil up to the level of the collar; the roots should not be exposed, and the stem should not be excessively buried in the soil. Various types of tools can be used for planting, including:

- motorised augers with a small diameter tip;
- manual augers, when working on existing vegetation, with the precaution of loosening the soil slightly more than necessary to contain the soil block.

In small areas, such as private gardens, normal gardening tools can also be used; even in this case, it is a good practice to loosen the soil a little more than is necessary to hold the underground part of the seedling (root ball or soil block).

Planting schemes may vary according to requirements and soil conditions. In all cases, **seedling must be planted in small, mono-specific groups** (see *BOX Even the eye wants its part!* at page 30).

The planting of seedlings in soil blocks is a particularly resource- and energy-intensive process, especially when large areas are to be covered, thereby requiring a significant number of seedlings. Furthermore, the establishment of young plants does not always yield the expected results. The success of this process is dependent on a number of factors that are not always predictable, including:

- the type of species and ecotype;
- the ability of the plant to adapt from the 'protected' conditions of the nursery to those encountered in the wild;
- the impact of unpredictable and potentially adverse climatic conditions after planting (such as prolonged drought or significant temperature fluctuations);
- trophic activity and uprooting by wild animals attracted to the young plants (especially crows, hares, and wild boars).

For the reasons mentioned above, if the goal is simply to enrich the composition of a 'hay-type' meadow, overseeding is suggested.

Please refer to Table 1 on page 72 for a list of the main species to be used.

Summary Sheet

Characteristics	perennial; prolonged and abundant flowering; high landscape value
Location	sunny areas; soils ranging from light to heavy, from well-drained to moist
Required skills	medium
Preliminary work	if work entails overseeding or planting seedlings in soil blocks, it is necessary to provide a short cut with the removal of the mowed material
Period of intervention	mid-October
Technique	restoration of abandoned meadows: a. recovery through mowing: implementation of appropriate management strategies, including 2(3) cuts during each growing season, with removal of mowed material b. recovery through species enrichment restoration of intensive meadows: a. recovery of regular mowing b. cessation of fertilisation c. species enrichment
Flowering period	April-October
Management	<ul style="list-style-type: none">• regular periodic mowing (2(3) cuts/year). Plan the first mowing in late spring/early summer, coinciding with the conclusion of the main grasses' maturation period (June)• plan more summer cuts in case of presence of invasive species
Remarks	<ul style="list-style-type: none">• use seeds or plants of native species sourced from certified nurseries and/or retailers

BOX: DO YOU HAVE A FIELD THAT HAS BEEN DORMANT FOR A WHILE? MAYBE IT WAS ONCE A PRODUCTIVE CROP LAND? WHY NOT TURN IT INTO A COLOURFUL FLOWERING MEADOW?

The objective of these interventions is the creation of meadows that are rich in species, with a particular focus on entomophilous species, and that are capable of supporting a diversified living community, with a specific emphasis on pollinating insects. The techniques employed necessitate a more substantial input in terms of locating the requisite plant material. In certain instances, the availability of particular equipment for its collection is also a prerequisite. Furthermore, these techniques necessitate a certain degree of experience and technical preparation.

Characteristics	perennial; prolonged and abundant flowering; high landscape value
Location	sunny areas; soils ranging from light to heavy, from well-drained to moist
Required skills	the establishment of a species-rich meadow requires a certain degree of expertise and technical proficiency in the sourcing and harvesting of plant material, as well as in the preliminary soil preparation work
Preliminary work	harrowing, false sowing, harrowing (or weeding)
Period of intervention	mid-October
Technique	three techniques: <ul style="list-style-type: none"> • by sowing a mixture of seeds of native species • by the use of species-rich hay • by brush-harvested seed
Flowering period	April-October
Management	<ul style="list-style-type: none"> • regular periodic mowing (2(3) mowings/year). Plan the first mowing in late spring/early summer, coinciding with the conclusion of the main grasses' maturation period (June) • consider additional summer mowing if invasive species are present

Remarks	<ul style="list-style-type: none">• identification of suitable donor meadows by qualified experts• if you decide to use hay or brush-harvested seed<ul style="list-style-type: none">- carefully assess the quality of the donor meadow and the degree of seed maturity- bear in mind that the material should be harvested around mid to late June, dried and stored until use• if you opt to sow seed mixes<ul style="list-style-type: none">- use seeds of guaranteed local origin for entomophilous species- use a low seeding density for grasses (max 10 g/sqm) and add 1-4 g/sqm of entomophilous species• in the case of mesophilous or meso-hygrophilous meadows, provide a light fertilisation with well-rotted manure, excluding chemical fertilisation• avoid fertilisation in the case of xeric, meso-xeric and wet meadows
----------------	--

If you want to learn more, you can consult the first volume of the series 'Guidelines for the creation and management of habitats for pollinator. I. AGRICULTURAL AREAS', which can be found at the following link: https://www.lifepollinaction.eu/?page_id=3806&lang=en

CREATION AND MANAGEMENT OF WILDFLOWER STRIPS

This sheet explains how to create and manage wildflower strips.

Flower strips are defined as strips of entomophilous herbaceous plants of varying width and linear shape created at the edge of a lawn in a municipal park or private garden, at the foot of a boundary wall, in an urban vegetable garden, at the edge of drains, canals and small watercourses.



Although they may vary, flower strips are generally modest in size, varying in length and ranging from 2-3 to 10 meters in width. They comprise a mixture of different native herbaceous plants: the general rule is that **wildflower strips should be rich in plant species that differ in size, height, floral morphology and flowering period.**

CREATION OF PERENNIAL FLOWER STRIPS: HOW TO DO IT

Perennial strips are **flower strips composed of plant species with a multi-year life cycle**. In many cases, the above-ground part of the plant dries up during the winter, but the root system continues to live, producing new shoots the following spring. The species composition varies in accordance with the environmental conditions of the site. In mesophilous conditions, the composition is analogous to that of meadows. Conversely, if the objective is to create a strip on a ditch or canal, hygrophilous species must be selected, that is, species adapted to living in environments with permanently water-rich soil. Perennial strips remain in place for several years, flowering annually. However, their maintenance necessitates the implementation of an appropriate management strategy.

1. Preliminary work

Preparatory work is limited and consists of mowing the existing grass to a height of between 3 and 5 cm. This is done with the aim of reducing competition between the existing species and those to be planted, thus creating a more favourable environment for their establishment.

2. The realisation

The creation of a strip involves the exclusive use of **seedlings in soil blocks**, i.e., plant material that has already developed. It is essential that the seedlings are planted after a hole has been drilled in the soil to a depth appropriate to the size of the underground part (root ball or soil block). The seedlings should be submerged in the soil up to the level of the collar, making sure that the roots are not exposed and that the stem is not



Seedlings in soil blocks produced in the nursery of Volpares (Palazzo dello Stella) from seeds of native species

excessively submerged. It is essential to use suitably developed seedlings (around the second year of life) with a soil block firmly attached to the roots and free of weeds. There are various types of tools that can be used for planting, but it is important to loosen the soil by a little more than is necessary to hold the soil block.

The planting scheme can vary depending on the requirements, but it is important that **the planting of seedlings occurs in small mono-specific groups** (see *BOX Even the eye wants its part!* on page 30). In the case of hygrophilous strips, it is also necessary to pay particular attention to the water fluctuation zone, which is defined as the area between the minimum and maximum water levels. In the sections of the bank in closer proximity to the water level, it is recommended that species capable of tolerating periods of submergence be situated.

In order to facilitate the management of the strip, it is recommended that late summer-flowering species be localised in designated areas, which should only be mowed at the end of summer (see below, Management and maintenance of wildflower strips).

In mesophilous flower strips, the transplanting of seedlings can be combined with the **sowing of a seed mixture** to increase species diversity and speed up the formation of the strip. To increase ground cover, the seed mix may also include seeds of less competitive grasses such as *Anthoxanthum odoratum*, *Briza media* and *Trisetaria flavescens*. It is recommended that the seeds are scattered between the seedlings at a target density of 4 g/sqm. The seed mixture should be no more than one third grass seed.

The **optimal period** for transplantation and potential sowing is **autumn**.

In the case of hygrophilous strips, if the outer edge of the strip facing the ditch or canal is colonised by tall weed species, it would be beneficial to prepare a seedbed and establish a permanent cover by sowing a mixture of commercial grasses that are suitable for wet soils. This approach would help to avoid the excessive proliferation, growth, and shading of the newly planted seedlings by the more vigorous weed species.

With regard to the principal species to be utilised, one may refer to the lists provided in Tables 1 and 2 on pages 72 and 78.

MANAGEMENT AND MAINTENANCE OF WILDFLOWER STRIPS

The **maintenance of perennial wildflower strips requires regular mowing**. In general, two cuts per year will maintain both species composition and richness. The first mowing should not take place before mid-June to allow species to flower and complete their vegetative cycle. The second mowing should take place between the end of August and the first half of September. In the case of the spread of invasive species such as Johnson grass (*Sorghum halepense*), the frequency and method of mowing may need to be changed, with an increase in the number of cuts if necessary. Mowing, especially in summer is particularly important in the first year/two years and should be repeated several times if the invasion is massive. It is also advisable to take precautions with regard to summer-flowering species. These should be confined to a limited number of clearly defined areas within the strip, as they will only require mowing on the second occasion, thus allowing them to complete their cycle. In the case of hygrophilous strips, where intervention is carried out along the banks of watercourses managed by Reclamation Consortia, there must be a prior agreement to keep at least the first 50 cm of the bank uncut throughout the flowering season, intervening only for autumn mowing.

The sequence of operations required to create a flower strip is illustrated below. The example concerns the creation of a flower strip at the edge of a municipal road, but the sequence of operations is also applicable to strips created in a park or a garden.



1. The preliminary operations for the establishment of a mesophilous strip on a roadside location entail the preparation of two planting rows;



2. The planting holes are then created along the rows. In the absence of an auger, or in the case of smaller areas such as private gardens, conventional gardening tools may also be employed as an alternative;



3. Transport and positioning of mesophilous herbaceous seedlings in soil blocks;



4. The seedlings are arranged in a regular, alternating pattern along the two planting rows at regular distances;



5. The planting of seedlings should be undertaken with due care and attention, ensuring that the collar of the seedlings is positioned at ground level.

Summary Sheet

Characteristics	perennial; moderate to abundant flowering; high landscape value
Location	sunny and sheltered areas; soils ranging from light to heavy, from well-drained to moist
Required skills	medium (in the case of mesophilous strips, the preparation of the soil and/or existing turf is necessary)
Preliminary work	cutting of the turf
Period of intervention	mid-October
Technique	manual transplantation in small mono-specific groups
Flowering period	May-September
Management	<ul style="list-style-type: none">• periodic mowing; schedule two mowings per year, with the first mowing after the first half of June• plan for multiple summer cuts in case of invasive species presence
Remarks	<ul style="list-style-type: none">• use plants of native species from certified nurseries and/or retailers• limit the first mowing to specific areas to allow summer-flowering species to complete their life cycle, subjecting them only to the second cut• plan the creation of wildflower strips well in advance, as native plants used for this purpose may not always be available from nurseries

CREATION AND MANAGEMENT OF SHRUB PATCHES AND FIELD HEDGES

This sheet illustrates the methods for the establishment and maintenance of shrub patches and hedges, within parks and gardens.

Shrub patches are formations that exhibit an areal (non-linear) shape, recreated on small marginal areas (typically between 50 and 100 sqm) and consist exclusively of shrub species. In contrast, **hedges** are linear formations, typically established along the edges of canals and ditches or within parks. In addition to shrubs, they may also comprise a tree component, with varying degrees of coverage. Shrub patches and hedges are important habitats for many animal and plant species providing essential resources and shelter. Many species also use them to move and spread within a hostile territory such as urban areas. A considerable number of native shrub species are highly attractive to numerous species of pollinating insects. Furthermore, the flowering of shrub species often occurs between late winter and late spring, thereby providing a source of sustenance for pollinating insects when the flowering resources provided by grassland species are not yet available.

BOX: EARLY FLOWERING SHRUB SPECIES

SPECIES	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
<i>Cornus mas</i>	Yellow											
<i>Cornus sanguinea</i>				White								
<i>Crataegus monogyna</i>			White									
<i>Ligustrum vulgare</i>			White									
<i>Prunus spinosa</i>	White											
<i>Rosa canina</i>			Pink									
<i>Viburnum lantana</i>			White									
<i>Viburnum opulus</i>				White								

LEGEND: The coloured squares represent the colour of the flowers and the flowering period.

In addition to providing suitable habitats for many wildlife species, these habitats offer numerous other services: hedges mitigate wind, improve the local microclimate, regulate run-off water and clean it of nutrient loads, stabilise canal banks, provide a trophic source for beekeeping (melliferous hedges), and produce small fruits (e.g., *Corylus avellana*, *Prunus* spp., *Sambucus nigra*, etc.).



Extensive early flowering ensures nutrient resources for pollinating insects when the flower resources provided by grassland species are not yet available



In addition to providing resources and shelter for many wildlife species, hedges improve the local microclimate, purify water from nutrient loads, stabilise canal banks, provide a trophic source for beekeeping (melliferous hedges) and produce small fruits

THE REALISATION OF SHRUB PATCHES AND FIELD HEDGES: HOW TO DO IT

The methods of intervention for the creation of hedges and shrub patches are similar; however, the species composition must be selected with consideration of the local soil characteristics⁸, in particular, moisture levels.

1. Soil preparation

The objective of preliminary soil preparation is to enhance soil functionality and quality while simultaneously reducing the incidence of spontaneous weed vegetation. Operations, to be carried out shortly before planting, should only affect the surface layers, with light milling limited to the planting strip. This can be achieved through the utilisation of manual equipment, such as a rotary tiller.

BOX: IF THE SELECTED AREA WAS UTILISED FOR AGRICULTURAL PURPOSES

The establishment of a hedge or shrub patch in a previously arable land can prove challenging due to the prior tillage and fertilisation practices that agricultural land has undergone. In such instances, it is necessary to implement a deeper tillage process in order to effectively disrupt any impermeable layers that frequently form between the typical tillage threshold and the underlying soil (approximately 30-50 cm). The presence of these layers can impede the optimal development of plant root systems, thereby jeopardising the success of the intervention. The procedure entails the implementation of deep ploughing to a depth of a minimum of 60-80 cm, which is then followed by the application of superficial milling and harrowing. It is then recommended that the following steps be undertaken:

1. deep tillage using a subsoiler;
2. superficial milling of the soil;
3. preparation of the planting area by harrowing.

If you want to learn more, you can consult the first volume of the series 'Guidelines for the creation and management of habitats for pollinator. I. AGRICULTURAL AREAS', which can be found at the following link: https://www.lifepollinaction.eu/?page_id=3806&lang=en

2. Establishment of the hedge and shrub patch

This intervention employs exclusively **young seedlings, aged between one and three years, in soil blocks**. The planting operations are conducted manually or with the assistance of manual equipment, including transplanters.

Prior to planting, a small hole should be excavated (with an average diameter of 10 cm and a length of 10 cm, dependent upon the dimensions of the soil block) into which the seedlings are placed. It is of the utmost importance to ensure that the plant's collar is positioned at ground level; the roots should never be exposed, nor should the stem be sunk too deep into the soil. The planting hole should then be filled with excavated soil, with slight pressure applied to the soil block to encourage root growth outside of the soil block itself.


⁸ Del Favero R., 1999. *Biodiversità e indicatori nei tipi forestali del Veneto, Regione del Veneto – Direzione regionale delle foreste e dell'economia montana e Accademia italiana di scienze forestali*, pp 335.

For the establishment of a **shrub patch**, the plants should be planted according to a **checkerboard grid pattern**, with the shrubs staggered between rows. The planting density varies depending on the size and shape that the plant will assume when mature. In the case of regular-shaped surfaces, typically square or circular, exceeding 250-300 sqm, it is recommended to plant **one plant every 2 m** along each row, with **rows spaced 3-3.5 m apart**. This results in a density of one plant every 7 sqm. In the case of regular-shaped areas of less than 250-300 sqm, or for larger but narrow and elongated surfaces, it is possible to reduce the distance between rows to 2 m, thus reaching a planting **density of one plant every 4 sqm**. In order to establish a **hedge**, it is essential to apply **mulch** after soil preparation. This limits competition from spontaneous vegetation on the young plants during the initial years following the intervention. The procedure entails the utilisation of **biodegradable materials, such as jute or coconut fibre sheets**, over the tilled soil prior to the planting of the seedlings. These materials must be firmly secured at the sides and heads of each continuous stretch of hedge. The seedlings are then planted at intervals corresponding to the incisions made in the sheets at the desired distances.


The distance typically used between seedlings is **1.5 m**. In cases where the tree component is also included, it is advised that a distance of **9 m** be maintained between each tree.

The species employed in the establishment of hedges and shrub patches are typically sourced from standard nursery production, thereby mitigating potential issues associated with sourcing. In the case of a hedge that incorporates a tree component, it is recommended that both the species selected and their position be considered with regard to the potential constraints and shading that they may exert once they have reached maturity.


Please refer to Tables 1 and 2 on page 72 and 78 for the main species to be used.



Planting schemes for hedges consisting exclusively of shrubs



Planting schemes for shrub patches



Planting schemes for mixed tree-shrub hedges

The images below illustrate the sequence of operations required to create a hedge or a shrub patch. The creation of shrub patches or hedges necessitates a certain degree of expertise, predominantly in the context of preliminary soil preparation. It is also crucial to adhere to the procedures depicted in Figures 3 and 4. Following soil preparation, mulching is essential to prevent competition from spontaneous vegetation and ensure the survival of young plants during their initial years after the intervention.



1. Surface preparation of the soil using a rotary tiller



2. Excavation of the hole using manual mechanical equipment or a motorised auger



3. Mulching of a circular area with a radius of at least 50 cm around the planting hole using biodegradable material such as coconut fibre or jute



4. Planting of seedlings inside the planting holes through an X-shaped slit in the mulch cover



5. Manual filling of the planting holes and soil compaction



6. Placement of the supporting pole



7. Localised fertilisation using organic-mineral fertilizer



MANAGEMENT AND MAINTENANCE OF HEDGES AND SHRUB PATCHES

In comparison to herbaceous habitats, shrub patches and hedges present **fewer management and maintenance challenges**. During the initial years following planting, it is crucial to ensure regular **mowing** to minimise competition from invasive herbaceous species. Once the seedlings have become established, the intensity and frequency of mowing can be reduced, and will only serve to ensure better development for the shrubs or trees. After the third year from planting, mowing of the invasive species may no longer be necessary.

Summary sheet	
Characteristics	perennial; abundant spring flowering; high landscape value
Location	sunny areas; soils ranging from light to heavy, from well-drained to moist
Required skills	medium (preparation of soil and/or existing turf required)
Preliminary work	soil preparation through light milling; operations should only affect the surface layers
Period of intervention	mid-October to mid-December
Technique	manual staggered planting for shrub patches and linear planting for hedges
Flowering period	January to July
Management	mowing of invasive herbaceous species during the first two years after planting
Remarks	<ul style="list-style-type: none">• use plants of native species from certified nurseries and/or retailers• for the creation of a hedge, apply mulch to limit competition from spontaneous vegetation using biodegradable jute or coconut fibre sheets



SECTION 4

GUIDE TO ASSESSING THE QUALITY OF PARKS, GARDENS AND COURTYARDS

The implementation of effective landscape management strategies in urban and suburban areas enables the harmonious coexistence of humans, plants, and animal species, including pollinators. It is feasible to create habitats in a variety of settings, including schoolyards, parks, gardens, urban vegetable gardens, flowerbeds, and roadsides, that can provide pollinators with the essential resources they require to sustain themselves, namely access to foraging areas, nesting sites, and protection during the winter months.

This section presents a straightforward questionnaire for the assessment of public (parks, schoolyards, flowerbeds) and private (gardens and vegetable gardens) green spaces, along with recommendations for readily implementable measures that can be taken to make these areas 'pollinator friendly'.

Instructions

The questionnaire is divided into three sections:

- 1) the first section contains general information about the site. While not essential, the inclusion of photographic material or a map of the area may prove beneficial in facilitating the assessment process;
- 2) the second part of the questionnaire allows for an evaluation of the foraging habitats present;
- 3) the third part of the questionnaire allows for an evaluation of the nesting habitats present.

Each section comprises a series of questions, arranged in sub-sections, for the assessment of particular elements.

Once the assessment is complete, the Checklist of actions to promote pollinators in yards, gardens, and parks can be utilised to identify the most appropriate actions to render areas more 'pollinator friendly'.

Site evaluation: calculate the score

1. assign a value in the 'Points' column (or 0 if not present) to each variable;
2. add up the scores and calculate a subtotal for each sub-section (e.g., 1a. Percentage of site composed of flowering plants);
3. calculate the total score for each section by summing up the scores of the sub-sections and copy these values into the summary table (e.g., Total Score Foraging Habitat (1a + 1b + 1c + 1d + 1e)) to obtain the overall assessment;
4. the scores obtained in the two sections will be employed to identify the aspects that are most lacking to understand the most appropriate interventions. For example, if the score for nesting sites is found to be low, actions that increase this score should take priority. Ideally, a good overall score should be at least 100 points.

GENERAL INFORMATION ON THE SITE

site name:	
owner/manager:	
address:	
define and describe the project area (public park, private garden, schoolyard, etc.)	

MAP OR DRAWING OF THE SITE, IF AVAILABLE	

SITE ASSESSMENT		
Assessment dates	BEFORE:	
	AFTER:	
Score your site	MAX	AFTER
Section 1: foraging habitat	100	
Section 2: nesting habitat	50	
TOTAL SCORE	150	

SECTION 1: FORAGING HABITAT

Pollinators require a diverse range of floral resources that are available throughout the growing season. Native flora is particularly effective in supporting a diverse array of pollinators and is crucial for the survival of numerous species.

1a. Percentage of the site containing entomophilous plants (excluding paved areas and buildings, as well as intensively mown lawn areas or areas with invasive or noxious species) (Figure 1). Entomophilous plants include native plants, ornamental flowers, flowering shrubs and trees.

Max score of 10



	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH  ON PAGE 69-71
Percentage cover of entomophilous plants. <i>Score 1 point for every 10% cover of entomophilous plant</i>	0 - 10			
subtotal 1a				



Figure 1: Examples of percentage cover of flowering species (all entomophilous plants have to be counted, even if they are not currently flowering)

1b. Percent of the site covered by native entomophilous plants (Figure 2).

Max score of 10

	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH  ON PAGE 69-71
<p>Percentage cover of native entomophilous plants. Score 1 point for every 10% cover of native entomophilous plants</p>	0 - 10			
<p>subtotal 1b</p>				





Figure 2: All three gardens have a high proportion of entomophilous plants. However, they differ in the proportion of native species

1c. Foraging habitats present on site (Figure 3).

Max score of 39


	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH PAGE 69-71
<p>Species-rich patches of the garden, presence of native wild plants. <i>Score 10 if any of these features are present. If they cover the whole yard/site, score 39 and move on to 1d</i></p>	10			
<p>Lawn with plants that flower even after mowing (e.g., clover, dandelion). Or mowing is less frequent than in the ordinary management approach.</p>	4			
<p>Areas with native entomophilous shrubs or trees (see Tables 1-2 for species list).</p>	10			
<p>Under the trees there are native entomophilous plants, rather than wood chips, turf or other exotic plants.</p>	5			
<p>Vegetable garden with plants that are pollinated by insects (e.g., apple trees, courgettes).</p>	5			
<p>Ornamental flowerbeds with annual or perennial plants that provide resources for pollinators, such as lavender, cornflower, chamomile.</p>	5			
subtotal 1c				



1d. Foraging habitats present at different times of the year.

**See Tables 1 and 2 on pages 72 and 78 for the flowering periods of some species.*







Max score of 30

	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH ON PAGE 69-71
SPRING-blooming species. Herbaceous plants, shrubs, or trees that flower between February and May. This includes fruit trees and some flowering weeds (e.g., dandelions), but does not include invasive species. <i>Score 1 point for each species (up to 10 points)</i>	0 - 10			
SUMMER-blooming. Herbaceous plants, shrubs, or trees that flower between June and August. <i>Score 1 point for each species (up to 10 points)</i>	0 - 10			
AUTUMN- blooming species. Herbaceous plants, shrubs, or trees that flower between September and November. <i>Score 1 point for each species (up to 10 points)</i>	0 - 10			
subtotal 1d				

1e. Resource-rich plants, plants for specialist insects and host plants for butterfly and moth caterpillars.

* See Tables 1 and 2 on page 72 and 78 for the list of species.

Max score of 11

	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH ON PAGES 72-78
<p>Resource-rich plants ( Table 1 and 2 on pages 72-78). Some native plants produce many resources that can support many species of pollinators. Score 0.5 point for each species (up to 5 points)</p>	0 - 5			  
<p>Plants for specialist insects ( Table 1 and 2 on pages 72-78). These plants provide pollen for specialist insects that only visit a narrow range of plants. Score 0.5 point for each species (up to 3 points)</p>	0 - 3			
<p>Plants for butterfly and moth caterpillars ( Table 1 and 2 on pages 72-78). To survive and grow to metamorphosis, the caterpillars of butterflies and moths need certain types of plants. Score 0.5 point for each species (up to 3 points)</p>	0 - 3			
<p>subtotal 1e</p>				


Total Score Foraging Habitats
(1a + 1b + 1c + 1d + 1e)

SECTION 2: NESTING & OVERWINTERING HABITAT

Pollinators nest and overwinter in a variety of places. About 70% nest on the ground, while 30% nest in cavities in dead wood, branches or stems. Bumblebee nests are often found under trees, tall grasses or hidden in the vegetation. Many insects, including some butterflies, moths, beetles, flies and bumblebee queens, overwinter under leaf litter or more generally in the ground. Having a variety of nesting features on your site will increase the likelihood of successful nesting and overwintering.

2a. Habitat for ground-nesting insects and other insects that seek shelter in the ground (Figure 4).

Max score of 25

	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH ON PAGES 69-71
<p>Presence of vegetated areas that are unowned (including shrubland and marginal areas). <i>Score 1 point for every 10% of unowned vegetated areas</i></p>	0 - 5			
<p>Areas with patchy vegetation or bare ground (areas with soil that is too compacted or disturbed or treated with pesticides are excluded). <i>Absent: 0; Present: 5</i></p>	0 - 5			
<p>Clump-forming native bunch grasses in the garden or in unowned areas <i>Absent: 0; Sparse: 2; Abundant: 5</i></p>	0 - 5			

<p>Presence of litter (dead leaves) from autumn to spring. Absent: 0; Sparse: 2; Abundant: 5</p>	<p>0 - 5</p>			
<p>subtotal 2a</p>				




Figure 4: Although considered 'unsightly', gardens with bare patches of soil provide ideal conditions for ground-nesting insects. Unlike ground wasps, many solitary bees are active for short periods, make very small nests and sting very rarely. In fact, you may not even be aware of their presence!

2b. Habitat for insects that nest in cavities or seek shelter in wood or stems (Figure 5).

Max score of 25


	SCORE	BEFORE	AFTER	HOW TO IMPROVE? CHECK POINTS WITH  ON PAGES 69-71
Presence of dead stems of flowering plants. <i>Absent: 0; Sparse: 2; Abundant: 5</i>	0 - 10			
Presence of shrubs. <i>Absent: 0; Sparse: 2; Abundant: 5</i>	0 - 5			
Presence of logs or dead trees. <i>Absent: 0; Sparse: 2; Abundant: 5</i>	0 - 5			
Brush piles or other woody debris. <i>Absent: 0; Sparse: 2; Abundant: 5</i>	0 - 5			
Rock piles or rock walls. <i>Absent: 0; Sparse: 2; Abundant: 5</i>	0 - 5			
subtotal 2b				




















Figure 5: Most cavity nesting species use existing cavities in dead trees, logs, shrubs or stems/stalks of large herbaceous plants. Bumblebees also nest under large clumps of grass. Stone walls or wooden fences may attract additional pollinator species

**Total Score Nesting & Overwintering Habitats
(2a + 2b)**












CHECKLIST OF ACTIONS

To Promote Pollinators in Yards, Gardens & Parks

ACTIONS IN PARKS AND GARDENS

Plant a native wildflower garden. Choose species that flower at different times of the year and are of great value to pollinators (check species with  in Table 1, page 72)	
Plant native tuft-forming species; in addition to providing a food source for butterflies, these plants are a nesting site for bees	
Turn most of the garden into a flowering habitat	
Plant native entomophilous species that flower in spring	
Plant spring-flowering tree or shrub species	
Plant native entomophilous species that flower in summer	
Plant summer-flowering tree or shrub species	
Plant native entomophilous species that flower in autumn	
Plant native trees that can provide food for butterflies and moths caterpillars (check species with  in Table 1, page 72)	
Plant species that provide food for specialist pollinator species (check species with  in Table 1, page 72)	
Gradually replace exotic or pollinator-unfriendly annuals and perennials with native entomophilous species	
If you have exotic plants in your garden, make sure they are not invasive and are attractive to pollinators	
Remove exotic species from your garden	
Ensure that no pesticides are used in or near your garden	

GARDEN MANAGEMENT

Avoid pesticides. Try to use less harmful alternatives	
Reduce mowing frequency and increase cutting height to allow species to flourish	
Leave trunks and dead branches	
Leave a thin layer of leaves in some areas of the garden to form litter	
Leave some patches of bare ground in the garden, but make sure it does not become too compacted	
Avoid plastic mulch or heavy wood chips, and treated wood chips	
Leave dead wildflower stems over winter; cut back them in early spring to provide new nesting sites	
Maintain areas with shrubs to increase nesting sites	
Leave some areas of the garden unmown to provide tall grass and increase nesting sites	
Add a water source for insects that need it to build their nests	
Seed a lawn with clover and other species that will continue to flower after mowing	

VEGETABLE GARDEN AND ORCHARD
















<p>Plant fruit trees and shrubs, using local varieties where possible, such as elderberries, blackberries and raspberries (check species in Table 2, page 78)</p>	
<p>Plant blackberry bushes that provide nesting sites for pollinators</p>	
<p>Plant herbs such as basil (<i>Ocimum</i>), mint (<i>Mentha</i>) and lavender (<i>Lavandula</i>)</p>	
<p>Plant vegetables and fruit trees that flower at different times</p>	
<p>Do not use pesticides in your vegetable garden or orchard. Consider more sustainable options</p>	

TABLE 1: SUPERFOOD & HOST PLANTS














Scientific name	Common name	Flowering period	Habitat	Type
<i>Achillea millefolium</i> L.	Common yarrow	May-Oct	Ge	
<i>Ajuga reptans</i> L.	Bugleweed	Dec-Jun	Ge	
<i>Allium angulosum</i> L.	Mouse garlic	May-Aug	H	
<i>Althaea officinalis</i> L.	Common marshmallow	Apr-Aug	H	
<i>Caltha palustris</i> L.	Marsh marigold	Feb-Jun	H	
<i>Campanula glomerata</i> L.	Clustered bellflower	May-Sep	Ge	
<i>Centaurea nigrescens</i> Willd.	Short fringed Knapweed	May-Dec	Ge	
<i>Cirsium arvense</i> (L.) Scop	Creeping thistle	Apr-Nov	Ge	
<i>Cirsium oleraceum</i> (L.) Scop.	Cabbage thistle	May-Sep	H	
<i>Cirsium palustre</i> (L.) Scop.	Marsh thistle	May-Sep	H	
<i>Daucus carota</i> L. subsp. <i>carota</i>	Wild carrot	Apr-Oct	X	
<i>Echium vulgare</i> L.	Blue viper's-bugloss	Mar-Sep	X	
<i>Eupatorium cannabinum</i> L. subsp. <i>cannabinum</i>	Hemp-agrimony	Jul-Oct	H	
<i>Genista tinctoria</i> L.	Dyer's Greenweed	Apr-Jul	H	

NATIVE WILDFLOWERS










SUPERFOOD & HOST PLANTS

NATIVE WILDFLOWERS	Scientific name	Common name	Flowering period	Habitat	Type
	<i>Gratiola officinalis</i> L.	Hedge hyssop	May-Aug	H	
	<i>Hypericum perforatum</i> L.	St. John's wort	Apr-Aug	Ge	
	<i>Inula britannica</i> L.	British yellowhead	Jun-Sep	H	
	<i>Iris pseudacorus</i> L.	Water flag	Mar-Jun	H	
	<i>Knautia arvensis</i> (L.) Coult.	Common ambretta	Apr-Sep	Ge	
	<i>Leontodon hispidus</i> L.	Bristly hawkbit	May-Oct	Ge	
	<i>Leucanthemum vulgare</i> Lam. subsp. <i>vulgare</i>	Ox-eye daisy	Feb-Oct	Ge	
	<i>Lysimachia vulgaris</i> L.	Yellow loosestrife	May-Aug	H	
	<i>Lythrum salicaria</i> L.	Purple loosestrife	May-Sep	H	
	<i>Malva alcea</i> L.	Greater musk-mallow	May-Sep	Ge	
	<i>Mentha longifolia</i> (L.) Huds.	Horsemint	May-Oct	H	
	<i>Onobrychis arenaria</i> (Kit.) DC.	Sand esparcet	Apr-Aug	X	
<i>Onobrychis viciifolia</i> Scop.	Sainfoin	Apr-Aug	Ge		
<i>Pastinaca sativa</i> L.	Wild parsnip	Jun-Aug	Ge		












SUPERFOOD & HOST PLANTS

NATIVE WILDFLOWERS	Scientific name	Common name	Flowering period	Habitat	Type
	<i>Salvia pratensis</i> L.	Meadow sage	Apr-Aug	X	
	<i>Scabiosa triandra</i> L.	Narrowleaf scabious	May-Sep	X	
	<i>Scrophularia umbrosa</i> Dumort. subsp. <i>umbrosa</i>	Green figwort	May-Sep	H	
	<i>Scutellaria galericulata</i> L.	Marsh Skullcap	May-Aug	H	
	<i>Senecio paludosus</i> L. subsp. <i>angustifolius</i> Holub	Fen Ragwort	May-Sep	H	
	<i>Silene flos-cuculi</i> (L.) Clairv.	Ragged-robin	Apr-Aug	H	
	<i>Stachys palustris</i> L.	Marsh Woundwort	May-Aug	H	
	<i>Symphytum officinale</i> L.	Comfrey	Apr-Nov	Ge	
	<i>Thalictrum lucidum</i> L.	Shining meadow-rue	May-Aug	H	
<i>Tragopogon pratensis</i> L. subsp. <i>orientalis</i> (L.) Celak.	Oriental goat's beard	May-Aug	Ge		
<i>Trifolium pratense</i> L.	Red Clover	Jan-Dec	Ge		
<i>Valeriana dioica</i> L.	Marsh Valerian	Apr-Jul	H		
<i>Veronica anagallis-aquatica</i> L. subsp. <i>anagallis-aquatica</i>	Water Speedwell	May-Oct	H		




SUPERFOOD & HOST PLANTS

NATIVE SHRUBS AND TREES	Scientific name	Common name	Flowering period	Habitat	Type
	<i>Acer campestre</i> L.	Hedge maple	Mar-May	Ge	
	<i>Carpinus betulus</i> L.	European hornbeam	Apr-Jun	Ge	
	<i>Cornus mas</i> L.	Cornel cherry	Jan-Apr	Ge	 
	<i>Cornus sanguinea</i> L.	European dogwood	Apr-Jun	Ge	 
	<i>Corylus avellana</i> L.	Hazelnut	Feb-Apr	Ge	 
	<i>Crataegus laevigata</i> (Poir.) DC.	Midland hawthorn	Mar-Jun	Ge	 
	<i>Crataegus monogyna</i> Jacq.	Common hawthorn	Mar-May	Ge	 
	<i>Euonymus europaeus</i> L.	Common spindle	Mar-Jun	Ge	
	<i>Frangula alnus</i> Mill. subsp. <i>alnus</i>	Alder buckthorn	Apr-Jun	H	 
<i>Fraxinus ornus</i> L. subsp. <i>ornus</i>	Flowering ash	Mar-May	Ge/X		
<i>Ligustrum vulgare</i> L.	Common privet	Mar-May	Ge		
<i>Lonicera caprifolium</i> L.	Perfoliate honeysuckle	Apr-Jul	Ge	 	
<i>Lonicera xylosteum</i> L.	European fly honeysuckle	Apr-Jul	Ge	 	

SUPERFOOD & HOST PLANTS

NATIVE SHRUBS AND TREES	Scientific name	Common name	Flowering period	Habitat	Type
	<i>Malus sylvestris</i> (L.) Mill.	Wild apple	Apr-May	Ge	
	<i>Prunus mahaleb</i> L.	Mahaleb cherry	Apr-May	Ge	
	<i>Prunus spinosa</i> L.	Blackthorn	Jan-Apr	Ge	
	<i>Pyrus communis</i> L. subsp. <i>pyrastrer</i> (L.) Ehrh.	Wild pear	Mar-May	Ge	
	<i>Rhamnus cathartica</i> L.	Common buckthorn	Mar-Jun	X	
	<i>Rosa canina</i> L.	Dog rose	Mar-Jul	Ge	
	<i>Salix cinerea</i> L.	Grey willow	Feb-Apr	H	
	<i>Salix purpurea</i> L.	Purple willow	Feb-Apr	H	
	<i>Sorbus torminalis</i> (L.) Crantz	Wild Service-tree	Mar-May	Ge	
	<i>Ulmus minor</i> Mill.	Field Elm	Feb-Mar	Ge	
	<i>Viburnum lantana</i> L.	Wayfaring Tree	Mar-May	H	
<i>Viburnum opulus</i> L.	Guelder-rose	Apr-Jun	Ge		

SUPERFOOD & HOST PLANT

	Scientific name	Common name	Flowering period	Habitat	Type
GRASSES	<i>Anthoxanthum odoratum</i> L.	Sweet vernal grass	Mar-Aug	Ge	
	<i>Briza media</i> L.	Quaking-grass	Apr-Aug	Ge	
	<i>Trisetaria flavescens</i> (L.) Baumg.	Yellow oatgrass	May-Aug	Ge	

Habitat: **Ge** = generalist; **H** = hygrophilous; **X** = drought tolerant (dry soil, dry grassland);



= 'SUPERFOOD' PLANTS - Some native plant species offer a substantial resource base for a diverse range of pollinators;

















= PLANTS FOR SPECIALIST INSECTS - Some insects are 'specialists', collecting pollen and nectar only on specific plants;








= PLANTS FOR BUTTERFLIES - The caterpillars of many butterfly species are very specialised and only feed on specific plants.

TABLE 2: EDIBLE PLANTS WITH VALUE TO POLLINATORS










Scientific name	Common name	Flowering period	Type
<i>Allium</i> sp.	garlic, chives, leeks, onions, shallots	various	
<i>Anethum graveolens</i>	dill	Jun-Aug	
<i>Brassica</i> sp.	broccoli, cauliflower	vari	
<i>Calendula officinalis</i>	marigold	May-Sep	
<i>Citrus</i> sp.	lemon, lime, mandarin, orange	various	
<i>Coriandrum sativum</i>	coriander	Apr-Giu	
<i>Cucumis</i> sp.	watermelon, melon	various	
<i>Cucurbita</i> sp.	pumpkin	various	
<i>Foeniculum vulgare</i>	fennel	May-Sep	
<i>Fragaria</i> sp.	strawberry	May-Jun	
<i>Helianthus annuus</i>	sunflower	Jun-Sep	
<i>Lavandula</i> sp.	lavender	May-Sep	
<i>Matricaria chamomilla</i>	chamomile	Apr-Aug	
<i>Mentha</i> sp.	mint	various	

HERBACEOUS SPECIES

EDIBLE PLANTS WITH VALUE TO POLLINATORS

HERBACEOUS SPECIES	Scientific name	Common name	Flowering period	Type
	<i>Ocimum basilicum</i>	basil	Apr-Sep	
	<i>Origanum vulgare</i>	oregano	May-Sep	
	<i>Phaseolus</i> sp.	bean	various	
	<i>Solanum</i> sp.	aubergine, potato, tomato	various	
	<i>Vicia</i> sp.	vetch	various	

EDIBLE PLANTS WITH VALUE TO POLLINATORS

Scientific name	Common name	Flowering period	Type
<i>Corylus avellana</i>	hazelnut	Feb-Apr	
<i>Castanea sativa</i>	chestnut	Apr-May	
<i>Malus domestica</i>	apple	Mar-May	
<i>Prunus</i> sp.	almond, apricot, cherry, peach	Mar-May	
<i>Pyrus communis</i>	pear	Mar-May	
<i>Rosa</i> sp.	rose	various	
<i>Rubus</i> sp.	blackberries, raspberries	Apr-Jun	
<i>Ribes</i> sp.	gooseberries	Apr-Jun	
<i>Sambucus nigra</i>	elderberry	Apr-Jun	

APPENDIX 1

INTERVENTION TIMELINE

	JAN		FEB		MAR		APR		MAY		JUN		JUL		AUG		SEP		OCT		NOV		DEC		
	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	1st	15th	
Meadow Improvement																									
Restoration of abandoned meadows																									
Option A. Recovery through mowing																									
1. Shrub clearing (if necessary)																									
2. Mowing with removal of clippings																									
Option B. Recovery through species enrichment																									
1. Mowing the existing turf to a height of between 3 and 5 cm																									
2a. Overseeding																									
2b. Enrichment through planting of native seedlings in soil blocks																									
Restoration of intensive meadows																									
1. Cessation of fertilisation practices																									
2. Mowing with removal of clippings																									
3. Species enrichment (see Option B)																									
Management and maintenance of the meadow																									
Mowing with removal of clippings																									

	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
	1-8	11-14	15-18	19-22	23-26	27-30	1-8	11-14	15-18	19-22	23-26	1-8
Establishment and Management of wildflower Strips												
Preliminary work												
1. mowing the existing turf to a height of between 3 and 5 cm												
Realisation of the strip												
Planting of native seedlings in soil blocks												
ONLY for mesophilous strips: seeding to support the planting												
Management and Maintenance of the Strip												
Mowing with removal of clippings												
Establishment and Management of Shrub Patches and Hedges												
Preliminary work												
Soil preparation through light milling												
Realisation of the hedge and shrub patch												
Planting of seedlings and mulching												
Management and maintenance of hedges and shrub patches												
Mowing of weed species in the first years												

NOTES:

A series of horizontal dotted lines for writing notes.

NOTES:

A series of horizontal dotted lines for taking notes.

NOTES:

A series of horizontal dotted lines for taking notes, starting below the 'NOTES:' header and extending across the width of the page.

NOTES:

A series of horizontal dotted lines for writing notes.



